REFERENCES

Abdalla, A., Klein, S., &Mäder, K. (2008). A new self-emulsifying drug delivery system (SEDDS) for poorly soluble drugs: characterization, dissolution, in vitro digestion and incorporation into solid pellets. European journal of pharmaceutical sciences, 35(5), 457-464.

Agarwal, V., Siddiqui, A., Ali, H., &Nazzal, S. (2009). Dissolution and powder flow characterization of solid self-emulsified drug delivery system (SEDDS). International journal of pharmaceutics, 366(1), 44-52.

Ajmera, A., Deshpande, S., Kharadi, S., Rathod, K., Patel, K., & Patel, P. (2012).Dissolution rate enhancement of atorvastatin, fenofibrate and ezetimibe by inclusion complex with β-cyclodextrin. Asian J Pharm Clin Res, 5(4), 73-76.

Anwar, M., Warsi, M. H., Mallick, N., Akhter, S., Gahoi, S., Jain, G. K., ...&Khar, R. K. (2011). Enhanced bioavailability of nano-sized chitosan–atorvastatin conjugate after oral administration to rats. European Journal of Pharmaceutical Sciences, 44(3), 241-249.

Aungst, B. J. (2000). Intestinal permeation enhancers.Journal of pharmaceutical sciences, 89(4), 429-442.

Balakrishnan, P., Lee, B. J., Oh, D. H., Kim, J. O., Hong, M. J., Jee, J. P., ...& Choi, H. G. (2009). Enhanced oral bioavailability of dexibuprofen by a novel solid self-emulsifying drug delivery system (SEDDS). European Journal of Pharmaceutics and Biopharmaceutics, 72(3), 539-545.

Bang, L. M., Chapman, T. M., & Goa, K. L. (2003). Lercanidipine. Drugs, 63(22), 2449-2472.

Bhagwat, D. A., & D'Souza, J. I. (2012). Formulation and evaluation of solid self micro emulsifying drug delivery system using aerosil 200 as solid carrier.International current pharmaceutical journal, 1(12), 414-419.

Chambin, O., Jannin, V., Champion, D., Chevalier, C., Rochat-Gonthier, M. H., &Pourcelot, Y. (2004). Influence of cryogenic grinding on properties of a self-emulsifying formulation. International journal of pharmaceutics, 278(1), 79-89.

Chen, Y., Li, G., Wu, X., Chen, Z., Hang, J., Qin, B., & Wang, R. (2008). Self-microemulsifying drug delivery system (SMEDDS) of vinpocetine: formulation development and in vivo assessment. Biological and Pharmaceutical Bulletin, 31(1), 118-125.

deSmidt, P. C., Campanero, M. A., &Trocóniz, I. F. (2004). Intestinal absorption of penclomedine from lipid vehicles in the conscious rat: contribution of emulsification versus digestibility. International journal of pharmaceutics, 270(1), 109-118.

Dedakia, J., &Majumdar, S. H. (2012). Formulation and Evaluation of Microemulsion Based Tablets of Acyclovir for Better Patient Compliance. International Journal of Pharmatech Research, 4, 1730-1734.

Dedhiya, M., Rastogi, S., &Chhettry, A. (2005).U.S. Patent Application No. 11/218,820.

Dixit, A. R., Rajput, S. J., & Patel, S. G. (2010).Preparation and bioavailability assessment of SMEDDS containing valsartan.AAPS pharmscitech, 11(1), 314-321.

El Maghraby, G. M., Osman, M. A., Abd-Elrahman, H. E., &Elsisi, A. E. (2014).Self emulsifyingLiquisolid tablets for enhanced oral bioavailability of repaglinide: In vitro and in vivo evaluation. Journal of Applied Pharmaceutical Science Vol, 4(09), 012-021.

Epstein, M. (2000).Lercanidipine: a novel dihydropyridine calcium-channel blocker. Heart disease (Hagerstown, Md.), 3(6), 398-407.

Grove, M., Müllertz, A., Nielsen, J. L., & Pedersen, G. P. (2006). Bioavailability of seocalcitol: II: development and characterisation of self-microemulsifying drug delivery systems (SMEDDS) for oral administration containing medium and long chain triglycerides. European journal of pharmaceutical sciences, 28(3), 233-242.

Guay, D. R. (2002). Update on fenofibrate. Cardiovascular drug reviews, 20(4), 281-302.

Gursoy, N., Garrigue, J. S., Razafindratsita, A., Lambert, G., & Benita, S. (2003). Excipient effects on in vitro cytotoxicity of a novel paclitaxel self-emulsifying drug delivery system. Journal of pharmaceutical sciences, 92(12), 2411-2418.

Gursoy, R. N., & Benita, S. (2004). Self-emulsifying drug delivery systems (SEDDS) for improved oral delivery of lipophilic drugs. Biomedicine & Pharmacotherapy, 58(3), 173-182.

Ha, E. S., Choo, G. H., Baek, I. H., Kim, J. S., Cho, W., Jung, Y. S., ... & Kim, M. S. (2015). Dissolution and bioavailability of lercanidipine–hydroxypropylmethylcellulose nanoparticles with surfactant. International journal of biological macromolecules, 72, 218-222.

Hauss, D. J. (Ed.). (2007). Oral lipid-based formulations: enhancing the bioavailability of poorly water-soluble drugs (Vol. 170). CRC Press.

Huan, D., Yi, T., Liu, Y., Xiao, L., & He, J. K. (2011). [Influence of silica on intestinal absorption of solid self-microemulsifying drug delivery systems]. Yao xuexuebao= ActapharmaceuticaSinica, 46(4), 466-471.

Humberstone, A. J., Porter, C. J., &Charman, W. N. (1996).A physicochemical basis for the effect of food on the absolute oral bioavailability of halofantrine.Journal of pharmaceutical sciences, 85(5), 525-529.

Jha, S. K., Karki, R., Venkatesh, D. P., &Geethalakshami, A. (2011).Formulation Development & Characterization of Microemulsion Drug delivery systems Containing Antiulcer drug.International journal of drug development and research.

Kang, B. K., Lee, J. S., Chon, S. K., Jeong, S. Y., Yuk, S. H., Khang, G., ... & Cho, S. H. (2004). Development of self-microemulsifying drug delivery systems (SMEDDS) for oral bioavailability enhancement of simvastatin in beagle dogs.International journal of pharmaceutics, 274(1), 65-73.

Kaushal, A. M., Gupta, P., &Bansal, A. K. (2004). Amorphous drug delivery systems: molecular aspects, design, and performance. Critical Reviews™ in Therapeutic Drug Carrier Systems, 21(3).

Kawakami, K. (2012). Modification of physicochemical characteristics of active pharmaceutical ingredients and application of supersaturatable dosage forms for improving bioavailability of poorly absorbed drugs. Advanced drug delivery reviews, 64(6), 480-495.

Kommuru, T. R., Gurley, B., Khan, M. A., & Reddy, I. K. (2001). Self-emulsifying drug delivery systems (SEDDS) of coenzyme Q 10: formulation development and bioavailability assessment. International journal of pharmaceutics, 212(2), 233-246.

Kumar, V. K., Devi, M. A., &Bhikshapath, D. V. R. N. (2013). Development of solid self emulsifying drug delivery system containing efavirenz: In vitro and In vivo evaluation. International Journal of Pharma and Bio Sciences, 4, 869-882.

Lawrence, M. J., & Rees, G. D. (2000).Microemulsion-based media as novel drug delivery systems. Advanced drug delivery reviews, 45(1), 89-121.

Leonetti, G., Magnani, B., Pessina, A. C., Rappelli, A., Trimarco, B., &Zanchetti, A. (2002).Tolerability of long-term treatment with lercanidipine versus amlodipine and lacidipine in elderly hypertensives*.American journal of hypertension, 15(11), 932-940.

Lindenberg, M., Kopp, S., &Dressman, J. B. (2004). Classification of orally administered drugs on the World Health Organization Model list of Essential Medicines ac-

cording to the biopharmaceutics classification system. European Journal of Pharmaceutics and Biopharmaceutics, 58(2), 265-278.

Lipinski, C. A. (2000). Drug-like properties and the causes of poor solubility and poor permeability.Journal of pharmacological and toxicological methods, 44(1), 235-249.

Mahmoud, E. A., Bendas, E. R., & Mohamed, M. I. (2009).Preparation and evaluation of self-nanoemulsifying tablets of carvedilol.AAPS pharmscitech, 10(1), 183-192.

Merisko-Liversidge, E., Liversidge, G. G., & Cooper, E. R. (2003).Nanosizing: a formulation approach for poorly-water-soluble compounds. European Journal of Pharmaceutical Sciences, 18(2), 113-120.

Merisko-Liversidge, E., Sarpotdar, P., Bruno, J., Hajj, S., Wei, L., Peltier, N., ...& Jones, J. (1996). Formulation and antitumor activity evaluation of nanocrystalline suspensions of poorly soluble anticancer drugs.Pharmaceutical research, 13(2), 272-278.

MHRA, Lercanidipine hydrochloride, Medicine and Healthcare products Regulatory Agency (2009) Public assessment report of lercanidipine hydrochloride 10 mg and 20 mg film coated tablets. 19

Nawale, R. B., & Mehta, B. N. (2013).Glibenclamide loaded self-microemulsifying drug delivery system (SMEDDS): development and optimization. Int J Pharm PharmSci, 5(2), 325-30.

Nazzal, S., & Khan, M. A. (2006).Controlled release of a self-emulsifying formulation from a tablet dosage form: Stability assessment and optimization of some processing parameters.International journal of pharmaceutics, 315(1), 110-121.

Nielsen, F. S., Gibault, E., Ljusberg-Wahren, H., Arleth, L., Pedersen, J. S., &Müllertz, A. (2007).Characterization of prototype self-nanoemulsifying formulations of lipophilic compounds.Journal of pharmaceutical sciences, 96(4), 876-892.

Nielsen, F. S., Petersen, K. B., &Müllertz, A. (2008). Bioavailability of probucol from lipid and surfactant based formulations in minipigs: influence of droplet size and dietary state. European Journal of Pharmaceutics and Biopharmaceutics, 69(2), 553-562.

October 2015. Available at: www.mhra.gov.uk/home/groups/par/documents/.../con134945.pdf

Pachava, S., Puttachari, S., sheriff, A., Thakur, RS. (2014). Formulation and Evaluation of Solid Self-microemulsifying Drug Delivery System of A Selective Second Generation Cephalosporin Antibiotic. Int. J. Pharm. Sci. Rev. Res., 24(2),176-181.

Patel, B. D., Modi, R. V., Thakkar, N. A., Patel, A. A., &Thakkar, P. H. (2012). Development and characterization of solid lipid nanoparticles for enhancement of oral bioavailability of Raloxifene.Journal of pharmacy &bioallied sciences, 4(Suppl 1), S14.

Pouton, C. W. (1997). Formulation of self-emulsifying drug delivery systems. Advanced Drug Delivery Reviews, 25(1), 47-58.

Pouton, C. W. (2000). Lipid formulations for oral administration of drugs: non-emulsifying, self-emulsifying and 'self-microemulsifying'drug delivery systems. European Journal of Pharmaceutical Sciences, 11, S93-S98.

Pouton, C. W. (2006). Formulation of poorly water-soluble drugs for oral administration: physicochemical and physiological issues and the lipid formulation classification system. European Journal of Pharmaceutical Sciences, 29(3), 278-287.

Pouton, C. W., &Charman, W. N. (1997).The potential of oily formulations for drug delivery to the gastro-intestinal tract.Advanced Drug Delivery Reviews, 25(1), 1-2.

Ranpise, N. S., Korabu, S. S., &Ghodake, V. N. (2014).Second generation lipid nanoparticles (NLC) as an oral drug carrier for delivery of lercanidipine hydrochloride. Colloids and Surfaces B: Biointerfaces, 116, 81-87.

Robinson, J. R. (1996). Introduction: Semi-Solid Formulations of Oral Drug Delivery. Bulletin Technique-Gattefosse, 11-14.

Schwendener, R. A., & Schott, H. (1996).Lipophilic 1-β-D-arabinofuranosyl cytosine derivatives in liposomal formulations for oral and parenteral antileukemic therapy in the murine L1210 leukemia model.Journal of cancer research and clinical oncology, 122(12), 723-726.

Serajuddin, A. (1999). Solid dispersion of poorly water-soluble drugs: early promises, subsequent problems, and recent breakthroughs. Journal of pharmaceutical sciences, 88(10), 1058-1066.

Sethia, S., &Squillante III, E. (2003). Solid dispersions: revival with greater possibilities and applications in oral drug delivery. Critical Reviews™ in Therapeutic Drug Carrier Systems, 20(2&3).

Shulman, M., Cohen, M., Soto-Gutierrez, A., Yagi, H., Wang, H., Goldwasser, J., ...&Nahmias, Y. (2011). Enhancement of naringenin bioavailability by complexation with hydroxypropoyl-β-cyclodextrin.PloS one, 6(4), e18033.

Tang, B., Cheng, G., Gu, J. C., &Xu, C. H. (2008). Development of solid self-emulsifying drug delivery systems: preparation techniques and dosage forms. Drug discovery today, 13(13), 606-612.

Thakkar, H., Nangesh, J., Parmar, M., & Patel, D. (2011).Formulation and characterization of lipid-based drug delivery system of raloxifene-microemulsion and self-microemulsifying drug delivery system.Journal of Pharmacy and Bioallied Sciences, 3(3), 442.

Tokumura, T., Tsushima, Y., Tatsuishi, K., Kayano, M., Machida, Y., & Nagai, T. (1987).Enhancement of the oral bioavailability of cinnarizine in oleic acid in beagle dogs.Journal of pharmaceutical sciences, 76(4), 286-288.

You, J., Li, Q. P., Han, X., Yu, Y. W., & Yang, M. S. (2005). A novel formulation design about water-insoluble oily drug: preparation of zedoary turmeric oil microspheres with self-emulsifying ability and evaluation in rabbits. International journal of pharmaceutics, 288(2), 315-323.